Phylogeography of commercial tuna and mackerel in the Indonesian Archipelago

¹ Department of Ecology and Evolutionary Biology, University of California Santa Cruz, 100 Shaffer Road, Santa Cruz, California 95060.

² Faculty of Fisheries and Marine Science, Diponegoro University, Semarang 50275, Indonesia. Email: <ambariyanto@undip. ac.id>.

³ Indonesia Marine Program JI, Conservation International, Drive Muwardi No. 17, Denpasar, Bali 80361, Indonesia. Email: <mverdmann@gmail.com>.

⁴ Faculty of Animal Science, Fisheries and Marine Science, The State University of Papua, Manokwari-West Papua 98314, Indonesia. Email: <hamid.toha@ yahoo.co.id>.

⁵ School of Medicine, Boston University, 72 East Concord Street, Boston, Massachusetts 02118. Email: <lukes87@gmail. com>.

⁶ Department of Ecology and Evolutionary Biology, University of California Los Angeles, 621 Charles E. Young Dr. South, Los Angeles, California 90095. Email: <paulbarber@ucla.edu>.

* Corresponding author email: <jackson@biology.ucsc.edu>.

Date Submitted: 10 December, 2012. Date Accepted: 4 July, 2013. Available Online: 3 December, 2013. Alexis M Jackson ^{1*} Ambariyanto ² Mark V Erdmann ³ Abdul Hamid A Toha ⁴ Luke A Stevens ⁵ Paul H Barber ⁶

ABSTRACT.—While numerous population genetics studies have investigated phylogeographic patterns of coral reef organisms in the Coral Triangle, few have addressed whether fishes in the pelagic environment exhibit concordant patterns of genetic subdivision. We analyzed approximately 400 base pairs of the mitochondrial control region to compare population structure and phylogeography of five pelagic tuna and mackerel within a subset of their geographic ranges (i.e., the Indonesian Archipelago). Focal species include frigate tuna [Auxis thazard (Lacépède, 1800)], kawakawa [Euthynnus affinis (Cantor, 1849)], skipjack tuna [Katsuwonus pelamis (Linnaeus, 1758)], Indian mackerel [Rastrelliger kanagurta (Cuvier, 1816)], and narrow-barred Spanish mackerel [Scomberomorus commerson (Lacépède, 1800)]. Observed patterns of regional genetic subdivision were consistent with the role of Pleistocene vicariance in structuring populations. Divergence dates of all pelagic fish lineages dated to the Pleistocene epoch. Concordant barriers to larval dispersal found near Sumatra, Sulawesi, and Papua suggested that the Halmahera and Mindanao eddies and the Indonesian flowthrough may be contemporary forces maintaining genetic divergence between demes of pelagic fishes. Given the economic importance of these species, we suggest that the scale of management for pelagics in Indonesia be re-evaluated to reflect regional differences in the genetic composition of fishes.

research paper

The Coral Triangle, which includes the marine environments of Indonesia, the Philippines, Papua New Guinea, the Solomon Islands, Malaysia, and East Timor, represents the global epicenter of marine biodiversity (Myers et al. 2000, Hughes et al. 2002, Roberts et al. 2002, Briggs and Bowen 2013). The exact processes generating patterns of increased species diversity in this region are unclear (Barber and

Bellwood 2005, Barber 2009, Bellwood and Meyer 2009), with multiple competing theories to explain the concentration of species diversity within this region of the world (e.g., Ekman 1953, Ladd 1960, Woodland 1983). However, vicariance is considered one of the key mechanisms driving population subdivision and lineage diversification in a wide variety of taxa within the Coral Triangle.

The Indonesian Archipelago represents a major biogeographic barrier between the Indian and Pacific Ocean provinces (Briggs 1974). During the Pleistocene, sea level fluctuations of up to 130 m exposed the Sunda and Sahul shelves, restricting waterways within the archipelago and isolating Indian and Pacific populations of marine species (Porter 1989, Voris 2000). Limited larval exchange between oceans, in conjunction with genetic drift and natural selection, facilitated genetic subdivision of demes. In addition to numerous biogeographic studies (e.g., McManus 1985, Woodland 1986, Springer and Williams 1990), a key line of evidence supporting Pleistocene vicariance between Indian and Pacific coral reef ecosystems comes from studies of genetic connectivity and phylogeography. While there are counterexamples in which Indo-Pacific phylogeographic structure is not observed in marine taxa (e.g., Klanten et al. 2007, Horne et al. 2008, Reece et al. 2010), there are numerous studies from the Coral Triangle that demonstrate largely concordant patterns of genetic structure in a wide variety of marine taxa (e.g., Williams and Benzie 1998, Benzie 1999, Lessios et al. 2001). Such concordant patterns may reflect signals of allopatric divergence from the Pleistocene (see Carpenter et al. 2011 and Barber et al. 2011 for review).

Genetics studies have also shown the role of ocean currents in reinforcing lineage divergence in the Coral Triangle. Ocean currents were once presumed to facilitate long distance dispersal and genetic homogeneity in marine systems (Sale 1991, Roberts 1997). Not only have marine populations proven more closed than initially expected (e.g., Jones et al. 1999, 2005, Almany et al. 2007), evidence has emerged to suggest that currents can inhibit larval dispersal and genetic connectivity. Barber et al. (2002) initially suggested the role of the Western Boundary Currents in limiting east to west larval transport across the Maluku Sea, a pattern later confirmed by Barber et al. (2006). Additional studies on mantis shrimp (Barber et al. 2011) as well as giant clams (Ravago-Gotanco et al. 2007, DeBoer et al. 2008, Kochzius and Nuryanto 2008, Nuryanto and Kochzius 2009), echinoderms (Crandall et al. 2008b), and fishes (Timm et al. 2008, Ackiss et al. 2013) have confirmed the role of currents in driving population subdivision. Recent biophysical modeling studies have also confirmed patterns of limited connectivity between Indo-Pacific demes (Kool et al. 2011).

The majority of phylogeography studies in the Coral Triangle have focused on either reef-associated or inshore taxa. To date few studies have investigated whether pelagic species encounter the same dispersal barriers as reef-dwelling organisms. Pelagic fishes are expected to exhibit weak population structure due to high dispersal potential, migratory behavior and large effective population sizes (Graves 1998, Ely et al. 2005). The few studies that have examined population structure of pelagic species in the Coral Triangle provide some evidence of east-west genetic subdivision among demes of pelagic fishes (Perrin and Borsa 2001, Rohfritsch and Borsa 2005, Sulaiman and Ovenden 2010). However, discordant geographic sampling across multiple unrelated studies makes direct comparison of these results challenging. As such, it remains unclear whether vicariance events and oceanographic processes that

Table 1. Biological data for focal species: Auxis thazard, Euthynnus affinis, Katsuwonus pelamis, Rastrelliger
kanagurta, and Scomberomorus commerson. Data include maximum length (max. length) in centimeters,
pelagic environment, depth range in meters, and spawning periodicity (Collete and Nauen 1983, Collette 1995,
Pauly et al. 1996, Riede 2004).

Species	Maximum length (cm)	Pelagic environment	Depth range (m)	Spawning periodicity
A. thazard	65	Neritic	50-?	Seasonal
E. affinis	100	Neritic	0-200	Seasonal
K. pelamis	110	Oceanic	0-260	Year round
R. kanagurta	35	Neritic	20-90	Seasonal
S. commerson	240	Neritic	10-70	Seasonal

promote population divergence in reef-dwelling species have subdivided the seemingly homogenous pelagic habitat in the same way as coral reef habitats.

We designed a phylogeographic study of five commercially important pelagic fishes. Focal species are members of the family Scombridae, including frigate tuna [Auxis thazard (Lacépède, 1800)], kawakawa [Euthynnus affinis (Cantor, 1849)], skipjack tuna [Katsuwonus pelamis (Linnaeus, 1758)], Indian mackerel [Rastrelliger kanagurta (Cuvier, 1816)], and narrow-barred Spanish mackerel [Scomberomorus commerson (Lacépède, 1800)]. While all five species are associated with pelagic environments (Collette and Nauen 1983), they vary in a number of biological characteristics including spawning periodicity, maximum size, and depth range (Table 1). We sampled within a subset of the entire geographic range for all species, focusing on the Indonesian Archipelago, a region well known for the presence of phylogeographic barriers and some of the world's highest landings of principal market tuna and mackerels (Majkowski 2007). We analyzed the hypervariable mitochondrial control region to investigate the role of both historical and contemporary processes such as Pleistocene sea level fluctuations and physical oceanography in generating regional patterns of lineage diversification within the archipelago. Knowledge of population structure and phylogeography may be used to inform fisheries management and conservation efforts for commercial pelagic species in Indonesia.

Methods

We collected tissue from the pectoral fins of *A. thazard* (n = 135, 11 localities), *E. affinis* (n = 188, 12 localities), *K. pelamis* (n = 177, 12 localities), *R. kanagurta* (n = 178, 11 localities), and *S. commerson* (n = 127, 9 localities) throughout the Indonesian Archipelago (Fig. 1, Table 2). Tissue samples were acquired in artisanal fish markets during 2005 and 2006 and preserved in 95% ethanol. Whenever possible, we collected all five species from the same markets to maximize the comparability of the datasets. Due to the small size of fishing vessels, lack of refrigeration and the high cost of petrol, artisanal fishermen only operate within waters 20–30 km from the fish markets. Local fishing activity reduces the possibility that fish sampled were shipped from other locations. We will consider these 20–30 km regions around fishing ports as separate sampling localities or demes in subsequent analyses.

We extracted DNA from tissue samples using a 10% Chelex (BioRad) solution (Walsh et al. 1991). We then amplified a 374–403 base pair fragment of control region using the primers CRA and CRE (Lee et al. 1995). Polymerase chain reaction (PCR) utilized the following thermocycler parameters: an initial hold at 94 °C for 5 min, 38 cycles of 94 °C for 30 s, 50 °C for 30 s, 72 °C for 60 s, followed by 72 °C for 7



Figure 1. Map of the Indonesian Archipelago showing sampling localities of pelagic tuna and mackerel (black circles). Numbers correspond to sample localities listed in Table 1.

min. PCR products were visualized on a 1% agarose gel. PCR products that amplified were cleaned by adding 5 ml of product to 5 units of Shrimp Alkaline Phosphatase (Biotech Pharmacon) and 5 units of Exonuclease I (GE Healthcare), incubating products at 37 °C for 30 min and 80 °C for 15 min. Cleaned PCR products were sequenced using BigDye terminator chemistry (Applied Biosystems, Inc.). We sequenced PCR products in the forward and reverse directions on an ABI 377 DNA sequencer. Sequences were proofread and aligned using Geneious v5.6 (Drummond et al. 2011).

We calculated molecular diversity and neutrality statistics for each species, including nucleotide diversity (π), haplotype diversity (h), Fu's F_s , Tajima's D, and Fu and Li's D*. Neutrality statistics can be used to uncover demographic phenomena and to detect selection in cases where patterns of DNA polymorphism deviate from those predicted by the Wright-Fisher neutral model of evolution (Fu and Li 1993, Tajima 1989, Fu 1996, Fu 1997). Simulation studies comparing the statistical power of such neutrality statistics revealed that F_s provides a powerful assessment of population expansion, while D* performs best when assessing background selection (Fu and Li 1993, Fu 1997). Following empirical work by Crandall et al. (2008a), we calculated these statistics to determine the relative importance of background selection and population expansions in demes of pelagic fishes.

We assessed phylogenetic relationships among sequences using haplotype networks and neighbor-joining trees, and plotted how patterns of genetic variation changed across geographic space. Haplotype networks based on pairwise distances between haplotypes were generated using Arlequin and R, and formatted using Adobe Illustrator. We generated neighbor-joining trees in PAUP (Swofford 1993). We estimated the average sequence divergence between clades, evaluating values >7% (McCune and Lovejoy 1998) for evidence of cryptic speciation or species misidentification. Finally, we plotted the frequency of distinct clades (with bootstrap support of 75% or higher) in each sampling locality as pie diagrams to illustrate the geographic distribution of genetic variation.

We tested for statistically significant population structure by estimating global and pairwise estimates of the fixation indices $F_{\rm ST}$ and $\Phi_{\rm ST}$, $F_{\rm ST}$ comes from foundational work by Sewall Wright looking at hierarchical partitioning of genetic variation among groups (Wright 1943, Wright 1951, Wright 1965). $\Phi_{\rm ST}$ is an analogue of $F_{\rm ST}$

rta,	
agu	
kan	
iger	
trell	н Э
Rasi	ity (
nis, .	/ersi
elan	e div
d sn	otido
пом	ucle
atsw), ni
s, Ka	s (n
ffini	lype
ns a	aplot
упп	of hê
Euth	ber (
rd, J	mn
haza	h), I
vis ti	ity (
Au	vers
cies	e di
spe	otyp
ocal	hapl
for f	(n),
ion	ens
reg	cim
ntrol	f spe
l coi	er ol
dria	nbe
hon	n, nı
nitoc	atio
he n	s loc
for t	nple
ces	. Sai
indi	uos.
sity	ıme
liver	соп
lar d	orus
lecu	rom
Мо	mbe.
le 2.	Sco
Tabi	and

		Auxis ti	hazard		E	uthynnus	affini	S	Kaı	nuomn;	s pela	nis	Rasi	relliger	kanag	urta	Scomb	eromoru	in com	nerson
Sample location	и	Ч	n_h	н	и	Ч	n_h	н	и	Ч	n_h	н	и	Ч	n_h	н	и	Ч	n_h	н
1. Aceh	13	1.00	13	0.031					22	1.00	20	0.063								
2. Medan	10	1.00	10	0.029	22	1.00	16	0.008	15	1.00	15	0.064	19	0.96	14	0.012	23	0.99	21	0.056
3. Nias					З	1.00	С	0.011	С	1.00	З	0.039					20	0.98	18	0.052
4. Padang																	С	1.00	С	0.012
5. Anyer					З	1.00	2	0.002												
6. Karimunjawa													5	1.00	5	0.008	23	1.00	22	0.048
7. Lombok	S	1.00	2	0.031																
8. Flores													С	0.00	1	0.000	20	0.96	15	0.030
9. Selayar	14	1.00	14	0.028	19	1.00	15	0.028												
10. Makassar	10	0.98	6	0.039	16	1.00	11	0.009	12	1.00	12	0.067	23	0.95	17	0.010	6	0.89	9	0.005
11. Manado	S	1.00	S	0.043					17	1.00	17	0.071	19	0.96	16	0.010				
12. Bitung					15	1.00	12	0.008	5	1.00	S	0.041								
13. Halmahera	27	1.00	27	0.032	18	1.00	13	0.009	20	1.00	20	0.064	20	0.94	15	0.011	13	1.00	13	0.052
14. Raja Ampat	15	1.00	15	0.042	16	1.00	11	0.031	19	1.00	19	0.059	19	0.95	13	0.010	10	1.00	10	0.035
15. Manokwari	15	0.99	14	0.035	18	1.00	13	0.058	20	1.00	20	0.065	19	06.0	11	0.016				
16. Biak	16	0.95	12	0.037	19	1.00	16	0.099	17	1.00	16	0.060	15	0.94	11	0.012				
17. Yapen	5	1.00	5	0.037	19	1.00	14	0.098	11	1.00	11	0.067	17	0.92	12	0.022	9	0.93	5	0.034
18. Jayapura					20	1.00	11	0.023	16	1.00	15	0.055	19	0.81	6	0.011				

that incorporates genetic distances between sequences to assess allelic correlations among groups (Excoffier et al. 1992). $\Phi_{\rm ST}$ is particularly informative in instances where $F_{\rm ST}$ is low as it less likely to underestimate population structure when allelic diversity is high (Hedrick 1999, Hedrick 2005, Bird et al. 2011). Statistical significance of pairwise estimates was assessed after Bonferroni correction (using a family-wise alpha rate of 0.05). We also assessed population structure of pelagics by calculating the standardized statistic $G''_{\rm ST}$, using unbiased estimates of heterozygosity within subpopulations ($H_{\rm S}$) and total heterozygosity ($H_{\rm T}$) generated in Arlequin (Meirmans and Hedrick 2011).

We then evaluated genetic differentiation across a previously reported region of vicariance in the archipelago, grouping sampling localities to reflect the separation of eastern and western Indonesia during Pleistocene low sea levels. Sampling localities were grouped into western (localities 1–8) and eastern Indonesia (sampling localities 9–18) (Fig. 1). We performed an Analysis of Molecular Variance (AMOVA) in Arlequin (Excoffier et al. 2005). Significance of AMOVAs was estimated using a permutation test with 10,000 replicates. Chi-squared analyses were performed concurrently using DnaSP v5.10 (Librado and Rozas 2009).

To determine the location of barriers to larval dispersal throughout region, we used the software Barrier (Manni et al. 2004). Barrier uses a computational geometry approach involving Delaunay triangulation and Voronoi tessellation to visualize patterns of genetic variation. Geographic distances are overlaid with genetic distances among sampling localities to detect the presence of barriers. We tested the significance of barriers by comparing geographic distance matrices against genetic distance matrices comprised of pairwise $F_{\rm ST}$ and $\Phi_{\rm ST}$ values estimated in Arlequin, performing 10 replicate analyses for each species. We will use the terms "strong" and "weak" barriers to indicate the relative frequency with which a given barrier appeared in the Barrier analyses. Strong barriers were confirmed by both $F_{\rm ST}$ and $\Phi_{\rm ST}$ pairwise estimates, while weak barriers were only seen for one fixation index or the other. We will use strength of barriers (depicted by their widths in Fig. 2) as a proxy for permeability of barriers and to identify potential phylogeographic breaks.

We empirically tested the role of Pleistocene vicariance events by generating coalescent genealogies. We calibrated a Bayesian skyline plot using a lognormal relaxed clock in BEAST v1.7.5 (Drummond et al. 2012). We allowed mutation rates to vary between 1% and 7% per million years (e.g., McCune and Lovejoy 1998) to generate lineage divergence dates. We used literature-derived mutation rates because fossils and calibration points were unavailable for our focal species. We used jModelTest 0.1.1 (Posada 2008) to select the nucleotide substitution model that best fit the control region data using Akaike information criterion. Analysis of sequences indicated that the GTR +I +G model best fit the data for *E. affinis, R. kanagurta,* and *S. commerson*, the GTR +G model best fit the data for *A. thazard* and the HKY +I +G model best fit the data for *K. pelamis.* To evaluate convergence of MCMC chains we ran each dataset three times for 50 million steps in BEAST. Effective sample size values of >200 suggested convergence as well as proper length and mixing of MCMC chains.

Finally, to test an alternative hypothesis to Pleistocene vicariance, we investigated whether isolation by distance could explain patterns of genetic subdivision in pelagic fishes. Limited dispersal across the archipelago (approximately 5000 km across) may result in a positive correlation between genetic distance and geographic distance. Pairwise Φ_{sT} values were generated in Arlequin and pairwise geographic distances



Figure 2. Pie diagrams for (A) Auxis thazard, (B) Euthynus affinis, (C) Katsuwonus pelamis, (D) Rastrelliger kanagurta, and (E) Scomberomorus commerson. Each pie chart illustrates the relative frequency of each clade among samples within a sampling locality. Dashed lines represent barriers to dispersal highlighted in BARRIER. Thick lines depict potential barriers that are supported by both F_{ST} and Φ_{ST} , while thin lines depict barriers that are only supported by significant values of one of these statistics.

between all sampling localities were estimated using a Geographic Distance Matrix Generator (Ersts 2013). Some geographic distances calculated in this way may include over land distances; however, this is not a major concern for this study as there are numerous potential pathways to measure distances between sampling sites on islands. Thus, such over land estimates may essentially average the difference between potential measured distances. We used Mantel and partial Mantel tests, implemented using the vegan package v2.0-9 in R (Oksanen et al. 2013), to test the statistical significance of isolation by distance. Partial Mantel tests assess the correlation between geographic distance matrices and genetic distance matrices while also controlling for the effect of hierarchal population structure. Hierarchal population structure can introduce bias to isolation by distance analyses (Meirmans 2012). Matrices for hierarchal population structure were coded with 0s and 1s to reflect whether pairs of populations were or were not in the same regional cluster. Partial Mantel tests were only performed in instances when we detected statistically significant hierarchal population structure in the AMOVA.

Results

We sequenced approximately 400 base pairs of control region for 805 tissue samples. Haplotype diversity was high across sampling localities for all five species, ranging from 0.81 to 1.00 (Table 2). For *A. thazard*, 100 sites were polymorphic, yielding 129 unique haplotypes from a total of 135 samples. For *E. affinis*, 143 sites were

polymorphic, yielding 109 unique haplotypes from a total of 188 samples. For *K. pelamis*, 165 sites were polymorphic, yielding 172 unique haplotypes from a total of 177 samples. For *R. kanagurta*, 36 sites were polymorphic, yielding 83 unique haplotypes from a total of 178 samples. For *S. commerson*, 103 sites were polymorphic, yielding 113 unique haplotypes from a total of 127 samples. Nucleotide diversity ranged from 0.002 to 0.099 between sampling localities for all five species (Table 2).

Significant departures from neutrality were observed in all five pelagics (Table 3). Tajima's D was negative and statistically significant for seven of 12 *E. affinis* sampling localities but was non-significant in the remaining four species. D* was also negative and statistically significant in five of 12 sampling localities of *E. affinis*. For the remaining four species, D* values were non-significant. In contrast, negative Fu's F_s values were statistically significant in all five species. Negative F_s and D* values indicate a departure from neutrality due to an excess of recent mutations (number of segregating sites greater than pairwise nucleotide diversity) from a potential population expansion or following a selective sweep. Although significantly positive values could result from sampling multiple distinct gene pools, for example as a result of an influx of divergent haplotypes from unsampled populations, our neutrality tests are uniformly negative. Given that samples for each locality were taken within the same 20 to 30 km region, they likely represent a single gene pool.

Neighbor-joining trees revealed one to two well-supported clades for each species, with no evidence found to suggest cryptic speciation (Online Fig. 1A–E). All samples for *K. pelamis* were considered a single clade due to low bootstrap support (<40%) for identified clades. Auxis thazard samples formed two clades, with 98% bootstrap support or higher, and 4.12% average sequence divergence between clades. Euthynnus affinis samples formed two clades, with 96% bootstrap support or higher, and 6.37% average sequence divergence between clades. Rastrelliger kanagurta formed two clades, with 79% bootstrap support or higher, and 2.15% average sequence divergence between clades. Scomberomorus commerson samples formed two clades, with 82% bootstrap support or higher, and 2.80% average sequence divergence averaged between clades. Visually, pie charts suggest regional trends in the distribution of clades among demes of A. thazard, E. affinis, R. kanagurta, and S. commerson within the Indonesian Archipelago. Multiple data sets confirm the presence of clades unique to eastern Indonesia [white clades for A. thazard (Fig. 2A), E. affinis (Fig. 2B), and *R. kanagurta* (Fig. 2D)]. In *S. commerson*, there is a decline in the abundance of the white clade from west to east across the archipelago (Fig. 2E). Pairwise Φ_{s_T} values reaffirmed these geographic trends (Online Tables 1A–E).

All five pelagics exhibited varying degrees of population structure across the Indonesian Archipelago (Table 4). We used previous categories defined by Wright (1965) to describe the magnitude of population structure observed in our focal species. Based on global Φ_{ST} values, we observed strong population structure in *R. kanagurta* ($\Phi_{ST} = 0.3728$, P < 0.0001) and *S. commerson* ($\Phi_{ST} = 0.3253$, P < 0.0001), moderate population structure in *A. thazard* ($\Phi_{ST} = 0.0953$, P < 0.0001) and *E. af-finis* ($\Phi_{ST} = 0.0173$, P = 0.0001) and weak population structure in *K. pelamis* ($\Phi_{ST} = 0.0113$, P = 0.0315). Based on F_{ST} values, we observed weak population structure in *A. thazard* ($F_{ST} = 0.0069$, P = 0.0001), *E. affinis* ($F_{ST} = 0.0074$, P < 0.0001), *K. pelamis* ($F_{ST} = 0.0313$, P = 0.0421), and *S. commerson* ($F_{ST} = 0.0225$, P < 0.0001), and moderate population structure in *R. kanagurta* ($F_{ST} = 0.0773$, P < 0.0001), and moderate population structure in *R. kanagurta* ($F_{ST} = 0.0773$, P < 0.0001), *G* "_{ST} values further supported genetic differentiation of populations for all five focal species. For *K.*

Table 3. Neutrality test statistics for the mitochondrial control region for focal species: Auxis thazard, Euthynnus affinis, Katsuwonus pelamis, Rastrelliger kanagurta and Scomberomorus commerson. Sample location, number of specimens (n), Tajima's D, Fu and Li's D* and Fu's F_s, as reported by Arlequin 3.11 and DnaSP v5.10.* denotes statistical significance of P < 0.05 for Tajima's D and Fu's F_s and P < 0.02 for Fu and Li's D*.

		Auxis	thazard			Euthymr	nus affini	s		Katsuwon	nus pelan	nis	Ru	astrellig	er kanag	urta	Scor	nberomo	rus comi	nerson
Sample location -	и	D	D*	$F_{\rm s}$	и	D	D*	F _s	и	D	D*	Fs	и	D	D*	Fs	и	D	D*	$F_{\rm s}$
1. Aceh	13	-1.02	-1.41	-5.55*					22	0.07	-0.26	-3.66								
2. Medan	10	-0.63	-1.42	-3.53*	22	-1.67*	2.48*	*79.9-	15	-0.21	-0.20	-3.61*	19	0.10	-0.15	-6.18*	23	-0.23	-0.15	-4.66*
3. Nias					ŝ	0.00		0.13	З	0.00		1.54					20	1.49	0.12	-3.42
4. Padang																	С	0.00		0.31
5. Anyer					ŝ	0.00		0.20												
6. Karimunjawa													5	-0.49	-0.19	-2.12*	23	-0.07	-0.12	-7.62*
7. Lombok	5	-0.39	-0.53	-3.76																
8. Flores													ŝ	0.00		0.00	20	-0.93	0.02	-2.29
9. Selayar	14	-1.46	-1.54	-6.72*	19	-2.28*	-3.64*	-3.02*												
10. Makassar	10	-0.43	-0.22	-1.01	16	-1.48	-2.19	-6.13	12	-0.03	-0.23	-2.18	23	0.03	-0.01	-8.53*	6	0.11	0.69	-2.33*
11. Manado	5	-0.01	-0.21	0.15					17	-0.27	-0.87	-4.20*	19	-0.02	-0.78	-9.24*				
12. Bitung					15	-1.67*	-1.65	-7.92*	5	-0.18	-0.18	0.21								
13. Halmahera	27	-1.20	-1.56	-9.59*	18	-1.58*	-2.01	-7.10*	20	-0.54	-0.58	-6.26*	20	-0.49	-0.29	-7.55*	13	0.00	0.22	-3.27*
14. Raja Ampat	15	-0.96	-1.27	-5.67*	16	-2.24*	-3.10*	-0.10	19	-0.20	-0.30	-6.10*	19	-0.13	-0.63	-5.70*	10	-0.64	-0.70	-2.70
15. Manokwari	15	-0.96	-1.08	-4.24*	18	-1.74*	-2.08*	0.93	20	-0.10	0.00	-6.26*	19	-0.30	0.28	-1.09				
16. Biak	16	-0.33	-0.52	-1.73	19	1.49	0.64	0.18	17	-0.37	-0.50	-2.87	15	-0.70	0.06	-3.46*				
17. Yapen	5	-0.37	-0.34	-0.04	19	-0.07	-0.68	2.14	11	-0.56	-0.79	-1.76	17	1.37	1.04	-1.57	9	-0.05	-0.31	1.28
18. Jayapura					20	-2.29*	-3.42*	0.27	16	-0.16	-0.08	-1.35	19	-1.21	-1.53	-0.61				

Table 4. Global estimates of $\Phi_{\rm ST}$ and $F_{\rm ST}$ for focal species: Auxis thazard, Euthynnus affinis, Katsuwonus pelamis, Rastrelliger kanagurta, and Scomberomorus commerson. * denotes statistical significance of P < 0.05 and ** of P < 0.001. P values were not provided for $G''_{\rm ST}$ since estimates were calculated manually.

Species	Φ_{st}	F _{ST}	$G''_{\rm ST}$
Auxis thazard	0.0980**	0.0069**	0.5843
Euthynnus affinis	0.1073**	0.0074**	0.1276
Katsuwonus pelamis	0.0113*	0.0313*	0.9995
Rastrelliger kanagurta	0.3728**	0.0773**	0.5356
Scomberomorus commerson	0.3253**	0.0225**	0.4599

pelamis, G''_{ST} approached a maximum value of one (Table 4). Results for *K. pelamis* in conjunction with the haplotype network suggest high levels of genetic differentiation are due to a lack of shared alleles among populations, with nearly every sequence representing a unique haplotype (Fig. 3C). G''_{ST} values for *A. thazard*, *E. affinis*, *R. kanagurta*, and *S. commerson* ranged between 0.1276 and 0.5843 (Table 4). Lower G''_{ST} values observed in the aforementioned species reflect both high haplotypic diversity and a higher proportion of shared alleles among demes (Fig. 3A,B,D,E).

Barriers to larval dispersal identified using Barrier are consistent with regional subdivision of groups of pelagic fishes (Fig. 2). Two strong barriers were identified in A. thazard separating western demes (Aceh, Nias, Lombok, Makassar, and Selayar) from two groups of eastern demes (Manado, Halmahera, Raja Ampat, and Manokwari; and Biak and Yapen). Weaker breaks were also observed between Aceh and Nias and between Halmahera and Raja Ampat. One strong barrier was observed in *E. affinis* separating Jayapura in Papua from the rest of the localities in the archipelago. Additional weaker barriers subdivided the archipelago into three groupings (Nias and Medan; Padang; and Makassar, Selayar, Bitung, Halmahera, Raja Ampat, Manokwari, Biak, and Yapen). Two strong barriers were observed in K. pelamis separating Aceh from a group of central (Nias, Medan, Bitung, and Manado) and eastern demes (Halmahera, Raja Ampat, Manokwari, Biak, Yapen, and Jayapura). Additional weaker barriers separated Bitung from Manado and Halmahera from Raja Ampat. Two strong barriers were observed in *R. kanagurta* subdividing the archipelago into three regional groups [western and central (Nias, Karimunjawa, Makassar, Manado, Halmahera, and Raja Ampat), southern (Flores), and eastern (Manokwari, Biak, Yapen, and Jayapura) Indonesia]. One strong barrier was observed in S. commerson, confirming the divide between western and eastern Indonesia observed in the AMOVA and chi-squared analysis. Additional weaker breaks were observed dividing Nias, Medan and Padang from one another.

We tested two potential hypotheses for regional subdivision of pelagic fishes: historical vicariance and isolation by distance. We found strong evidence to support the role of Pleistocene vicariance in structuring demes of pelagics. AMOVAs supported partitioning of genetic variation between eastern and western Indonesia in *S. commerson* ($\Phi_{CT} = 0.113$, P = 0.0445), with 11.31% of variation between regions, 16.62% within regions, and 72.07% within populations (Table 5). A chi-squared analysis further supported this trend ($\chi^2 = 123.257$, P = 0.0177, df = 105). Coalescent analyses revealed that all individuals sampled for each species coalesced to a most recent common ancestor that dated within the Pleistocene epoch (approximately 2,500,000 to 11,800 years ago) (Fig. 4). Results are inconsistent with the notion that observed patterns of genetic subdivision of pelagic demes reflect an ancestral divergence of 5–20



Figure 3. Haplotype networks for (A) *Auxis thazard*, (B) *Euthynnus affinis*, (C) *Katsuwonus pelamis*, (D) *Rastrelliger kanagurta*, and (E) *Scomberomorus commerson*. Circles are sized proportionally to the number of individuals that possess each haplotype. All haplotypes are separated by one mutation step unless denoted with hatch marks or a number. Haplotypes are color-coded based on associated clades in neighbor-joining trees.

Table 5. AMOVA results for focal species, *Auxis thazard, Euthynnus affinis, Katsuwonus pelamis, Rastrelliger kanagurta,* and *Scomberomorus commerson,* showing degrees of freedom (df), variance components (var), percent variation (% var) and *F* statistics to test regional hypothesis for vicariance between western and eastern Indonesia. * denotes statistical significance of P < 0.05. Φ_{CT} represents the correlation of random haplotypes within a group of populations in a region relative to that of random pairs drawn from the entire species. Φ_{SC} represents the correlation of random haplotypes within populations relative to random pairs of haplotypes in the entire species.

		Am	ong regi	ons	Amon	g popula	ations wit	hin regions		Within	populati	ons
Species	df	var	% var	$\Phi_{\rm CT}$	df	var	% var	Φ_{sc}	df	var	% var	Φ_{st}
A. thazard	1	0.02	0.28	0.003	9	0.59	9.50	0.095*	134	5.64	90.22	0.098*
E. affinis	1	-1.70	-0.14	-0.017	10	0.99	12.43	0.122*	176	7.11	89.27	0.107*
K. pelamis	1	-0.01	-0.08	-0.001	10	0.15	1.21	0.012*	165	11.86	98.87	0.011*
R. kanagurta	1	-0.15	-3.93	-0.039	9	1.57	41.22	0.397*	167	2.39	62.72	0.373*
S. commerson	1	1.75	14.66	0.147*	7	2.14	17.86	0.209*	118	8.07	67.47	0.325*



Figure 4. Coalescent genealogies generated in BEAST showing divergence dates for (A) *Auxis thazard*, (B) *Euthynnus affinis*, (C) *Katsuwonus pelamis*, (D) *Rastrelliger kanagurta*, and (E) *Scomberomorus commerson*. Divergence dates are listed in years.

Myrs. Within clades, the majority of individuals began to diverge between 326,000 and 64,000 years ago; 68,000 and 15,000 years ago for *E. affinis*; 470,000 and 400,000 years ago for *K. pelamis*; 54,000 and 19,000 years ago for *R. kanagurta*; and 120,000 and 41,000 years ago for *S. commerson. Euthynnus affinis* and *R. kanagurta* explicitly show some lineages that have diverged since the last glacial maximum (26,000 to 19,000 years ago). In contrast, we found no support for isolation by distance. All Mantel tests had non-significant correlation coefficients for *A. thazard* (r = -0.1355, P = 0.7683), *E. affinis* (r = -0.1065, P = 0.7588), *K. pelamis* (r = 0.0070, P = 0.4111), *R. kanagurta* (r = -0.0251, P = 0.4402), and *S. commerson* (r = 0.1050, P = 0.2309). We performed a single partial Mantel test due to an instance of hierarchal population structure in *S. commerson*. The partial Mantel test also yielded non-significant correlation coefficients after controlling for both geographic distance (r = 0.0966, P = 0.2720) and hierarchal population structure (r = 0.0773, P = 0.2697).

DISCUSSION

Pelagic fishes are expected to exhibit weak genetic differentiation over large geographic distances due to their large effective population sizes and potential for long distance dispersal (Ward et al. 1994, McOuinn 1997, Nesbo et al. 2000). Numerous empirical studies have shown an absence of genetic structure in pelagic fishes (Graves et al. 1984, Elv et al. 2005, Santos et al. 2010, Kumar et al. 2012). In instances where strong genetic subdivision was observed it was seen between demes in separate ocean basins (Chow and Ushiama 1995, Bremer et al. 2005, Lu et al. 2006, Diaz-Jaimes et al. 2010). Limited studies of pelagics within the Indonesian Archipelago have been unable to find evidence of genetic structure in nearshore or epipelagic fish species (e.g., Arnaud et al. 1999, Santos et al. 2010). In contrast, our study of five pelagic fishes shows strong evidence of regional genetic differentiation across the Indonesian Archipelago. Genetic structure ranged from a minimum of $\Phi_{sT} = 0.011$ ($F_{sT} = 0.0313$) in *K. pelamis* up to $\Phi_{sr} = 0.373$ ($F_{sr} = 0.0773$) in *R. kanagurta*, showing high variation among haplotypes and strong regional differences in haplotype frequencies. Both historical and contemporary processes have likely limited genetic exchange within Indonesia, subdividing demes of pelagics throughout the archipelago.

Evidence of Barriers to Dispersal in Indonesian Pelagic Fishes

Numerous phylogeographic breaks observed in marine taxa in the Coral Triangle have been attributed to genetic divergence driven by Pleistocene sea level fluctuations (McMillan and Palumbi 1995, Williams and Benzie 1998, Benzie 1999, Barber et al. 2000, Carpenter et al. 2011). Barriers to larval dispersal observed in our datasets are similar to those seen in both invertebrate and fish species within the Indonesian Archipelago. Concordance of some barriers suggest similar responses of marine organisms to past vicariance events. Variation observed in the exact placement of some barriers may indicate the presence of tension zones, which shift in space based on the balance between spatially variant selection and dispersal (Barton and Hewitt 1985). Despite these minor variations in placement, the majority of phylogeographic barriers observed could be categorized in the following ways: barriers isolating western Indonesian localities near Sumatra, barriers isolating localities in Papua and broad scale regional barriers isolating eastern and western Indonesia.

Divergence in Western Indonesia.—The first major category of barriers were observed in *A. thazard, K. pelamis, R. kanagurta,* and *S. commerson.* Similar phylogeographic breaks have been observed in both sessile and motile invertebrates (Barber et al. 2002, 2006, DeBoer et al. 2008, Kochzius and Nuryanto 2008, Vogler et al. 2008), as well as in damselfish, sygnathids, and caesionids (Lourie et al. 2005, Timm et al. 2008, Drew and Barber 2009, Leray et al. 2010, Ackiss et al. 2013). Divergence of western Indonesian demes including Medan, Nias, and Aceh has likely resulted from reduced pelagic marine habitats, exposure of the Sunda Shelf during Pleistocene low sea stands, as well as the cooling of surface water temperatures facilitated by upwelling (Fleminger 1986, Springer and Williams 1990). Combined, these processes may have led to the extirpation of marine populations across much of the archipelago and drastically reduced adult and larval dispersal. Additionally, since Sumatra represents one of the westernmost boundaries of the Indo-Pacific barrier, we might predict evidence of increased intraspecific isolation among demes in this region.

Divergence in Eastern Indonesia.—A second major category of barriers were observed in *A. thazard, E. affinis, K. pelamis,* and *R. kanagurta.* Demes in eastern Indonesia were genetically divergent, particularly those in Teluk Cenderawasih such as Manokwari, Biak, and Yapen. These findings contribute to a growing body of literature indicating genetic divergence in this region for multiple reef-dwelling invertebrates and some fishes (Barber et al. 2006, 2011, Crandall et al. 2008b, DeBoer et al. 2008, Timm et al. 2008, Nuryanto and Kochzius 2009). The large number of recently described endemic reef fishes (e.g., Allen and Adrim 2000, Allen and Erdmann 2006, Allen 2008, Gill and Allen 2011) further support the patterns of geographic isolation observed in marine species in eastern Indonesia.

Genetic divergence in eastern Indonesia could be driven by three mechanisms. Barber et al. (2002, 2011) suggested that the Halmahera Eddy may facilitate divergence. The Halmahera Eddy is an anticyclonic gyre that forms south of the North Equatorial Countercurrent (Wvrtki 1961, Arruda and Nof 2003). The eddy may be responsible for larval retention within Teluk Cenderawasih (includes Biak, Manokwari and Yapen), with water masses moving in a closed loop at speeds of up to 50 cm s^{-1} (Lukas et al. 1991) potentially limiting east to west larval transport between demes of pelagic fishes. The Mindanao Eddy may also drive genetic divergence among groups depending on the monsoon season (Schiller et al. 2008). The presence of a sill at 10–200 m in depth (Hall 2002) across Teluk Cenderawasih is a second mechanism that may facilitate genetic divergence in eastern Indonesia. During Pleistocene low sea levels stands, the exposure of this sill could have constricted water flow and larval dispersal in and out of Teluk Cenderawasih, promoting vicariance and genetic divergence in the region. Finally, it has been hypothesized that a combination of tectonic activity and changing sea levels has resulted in numerous isolations of Teluk Cenderawasih over the past 5 Myrs (Allen and Erdmann 2006).

Divergence Near the Makassar Strait.-The final major category of barriers are observed in the vicinity of Sulawesi at the Makassar strait in S. commerson and E. affinis, and the Maluku Sea in A. thazard and K. pelamis. These barriers roughly divide the archipelago into eastern and western Indonesia. An observed barrier at the Maluku Sea mirrors a historical biogeographic break known as Weber's Line (van den Bergh et al. 2001). Phylogeographic breaks at Weber's Line have been seen in mantis shrimp, nautilus, and sygnathids (Wray et al. 1995, Lourie et al. 2005, Barber et al. 2011). A break at the Makassar Strait has been seen in a few reef-dwelling and pelagic fish species (Perrin and Borsa 2001, Lourie et al. 2005). The Makassar Strait represents the primary passageway of the Indonesian Throughflow, which dominates the oceanography of the Indonesian Archipelago, and represents a dispersal pathway bringing water from the Pacific Ocean to the Indian Ocean (Godfrey 1996, Gordon and Fine 1996). The presence of a barrier to larval dispersal for pelagic fishes at the Makassar strait may reflect inter-decadal variability in water pathways due to El Nino Southern Oscillations (Valsala et al. 2011), thereby restricting regional connectivity and reinforcing historical genetic divergence among demes.

Demographic History

Our genetic data sets provide strong evidence to support a shared demographic history among pelagic fishes in the Indonesian Archipelago. Statistically significant Fu's F_s values in all five species, in conjunction with star polytomies observed in haplotype networks and coalescent analyses, indicate populations of pelagic fishes have

either recovered from a selective sweep or undergone a recent population expansion. Potential signals of population expansion are likely a signature of rising sea levels after the Pleistocene. Species once occupying the Sunda and Sahul shelves would have experienced a series of local extinctions, followed by recolonization events as sea levels rose. Inter-specific differences were, however, observed in Fu and Li's D*. Statistically significant values were only observed in one of the tuna species, *E. affinis*. Values indicate that strong population structure observed in *E. affinis* may partly result from background selection.

Relevance To Conservation And Fisheries Management

Genetic analyses are potentially invaluable to management of commercial species (Utter 1991, Carvalho and Hauser 1994). Failure to properly delineate and manage distinct fish stocks can result in local overfishing, species declines and a lack of recovery in exploited stocks (Cook et al. 1997, Hutchings 2000). Indonesia has significant economic incentives for protecting its marine resources given that marine fishing activity employs about 3.3 million people and annual exports of *K. pelamis* and other tunas approximate \$68 million USD (FAO 2004, Hutagalung 2012). Results from our study reinforce management unit boundaries proposed within the archipelago in Carpenter et al. (2011), with geographic isolation observed in demes in Sumatra and Teluk Cenderawasih. Additionally, the recent shift towards more local scale management of marine resources in Indonesia is favorable given evidence of regional subdivision of pelagic species. Localized fisheries management plans focus on implementation of community-based management initiatives (Christie 2005, Ablan 2006) and more spatially explicit strategies (Planes et al. 2009).

Our findings also allow us to make inferences about which regions in the archipelago may require more immediate management and conservation action. Regional genetic subdivision of demes in western and eastern Indonesia aligns with regional differences in fishing pressure. Ports with the highest annual landings are found in western Indonesia (FAO 2004, Ministry of Marine Affairs and Fisheries 2011), with the majority of all fishers operating from the islands of Sumatra and Java (Dwipongoo 1987, Cardinale et al. 2009). While evidence suggests that stocks in eastern Indonesia are less exploited, in recent years this region has experienced the greatest increase in landings (FAO 2004, Ministry of Marine Affairs and Fisheries 2011). Given observed genetic divergence of demes off western Sumatra and high levels of exploitation, commercial species surrounding Sumatra may require more immediate management attention.

Limitations of Single Locus Study

Finally, we would like to briefly address potential limitations of inferences made from our results given the use of a single mitochondrial marker. Analysis of a single molecular marker may only reveal a small portion of the evolutionary history of a species and in some instances provide unreliable assessments of genome-wide heterozygosity. Despite the limitations, our study provides evidence of multiple species displaying similar responses to evolutionary processes and biogeographic events. The Coral Triangle represents a well-studied region in which a number of concordant phylogeographic patterns have been observed across a wide variety of marine taxa. A large portion of the seminal work in this region was based on analysis of single locus or mitochondrial DNA data sets (i.e., McMillan and Palumbi 1995, Lavery et al. 1996, Palumbi et al. 1997), with analysis of additional markers often times reconfirming phylogeographic patterns. Our study provides an excellent starting point for understanding population dynamics of this under-studied group of fishes across the Indonesian Archipelago.

In conclusion, using the mitochondrial control region, we investigated population structure and phylogeography of five pelagic fishes in the Indonesian Archipelago. Our study revealed population structure in all five species and evidence of divergent populations across multiple taxa in both western Sumatra and Papua. Regional patterns of genetic subdivision in pelagic fishes are highly consistent with the timing of Pleistocene vicariance events, as seen for numerous other marine species in the Indonesian Archipelago. Future work could involve simulation studies to explicitly test whether lineage diversification can be directly linked to the presence of observed phylogeographic barriers. Nonetheless, our improved knowledge of regional subdivision of tuna and mackerel species provides crucial information to enhance management of commercial pelagic fishes in the Indonesian Archipelago.

Acknowledgments

We would like to acknowledge the Indonesian government and the Indonesian Institute of Sciences (LIPI) for collections permits (1187/SU/KS/2006 and 04239/SU.3/KS/2006). JK Barber-Choi, J Drew, E Crandall, S Pardede, T DeBoer, C Starger, K Chavarria, M Subia, M Muñoz, A Bradshaw, and M Sindhu assisted with field collections and logistics. E Crandall and G Bernardi provided helpful comments on earlier versions of the manuscript. We would also like to acknowledge significant feedback provided from four anonymous reviewers. Funding was partially provided by the National Science Foundation (Biological Oceanography, OCE-0349177, and Population Biology, DEB-0338566) as part of "The Diversity Project." Additional funding was provided by the David and Lucile Packard Foundation's "Ecosystem Based Management of the Bird's Head Seascape" project and the Walton Family Foundation's Bird's Head Seascape initiative.

LITERATURE CITED

- Ablan MCA. 2006. Genetics and the study of fisheries connectivity in Asian developing countries. Fish Res. 78(2–3):158–168. http://dx.doi.org/10.1016/j.fishres.2006.02.006
- Ackiss AS, Pardede S, Crandall ED, Ablan-Lagman M, Ambariyanto, Romena N, Barber PH, Carpenter KE. 2013. Pronounced genetic structure in a highly mobile coral reef fish, *Caesio cuning*, in the Coral Triangle. Mar Ecol Prog Ser. 480:185–197. http://dx.doi.org/10.3354/meps10199
- Allen GR. 2008. Conservation hotspots of biodiversity and endemism for Indo-Pacific coral reef fishes. Aquat Conserv. 18(5):541–556. http://dx.doi.org/10.1002/aqc.880
- Allen GR, Adrim M. 2000. *Amblypomacentrus clarus*, a new species of damselfish (Pomacentridae) from the Banggai Islands, Indonesia. Rec Aust Mus. 20(1):51–55.
- Allen GR, Erdmann MV. 2006. *Cirrhilabrus cenderawasih,* a new wrasse (Pisces: Labridae) from Papua, Indonesia. Aqua. 11:125–131.
- Almany GR, Berumen ML, Thorrold SR, Planes S, Jones GP. 2007. Local replenishment of coral reef fish populations in a marine reserve. Science. 316(5825):742–744. PMid:17478720. http://dx.doi.org/10.1126/science.1140597
- Arnaud S, Bonhomme F, Borsa P. 1999. Mitochondrial DNA analysis of the genetic relationships among populations of scad mackerel (*Decapterus macarellus, D-macrosoma,* and *D. russelli*) in South-East Asia. Mar Biol. 135(4):699–707. http://dx.doi.org/10.1007/s002270050671

- Arruda WZ, Nof D. 2003. The Mindanao and Halmahera eddies—twin eddies induced by nonlinearities. J Phys Oceanogr. 33(12):2815–2830. http://dx.doi. org/10.1175/1520-0485(2003)033<2815:TMAHEE>2.0.CO;2
- Barber PH. 2009. The challenge of understanding the Coral Triangle biodiversity hotspot. J Biogeogr. 36(10):1845–1846. http://dx.doi.org/10.1111/j.1365-2699.2009.02198.x
- Barber PH, Bellwood DR. 2005. Biodiversity hotspots: evolutionary origins of biodiversity in wrasses (Halichoeres : Labridae) in the Indo-Pacific and new world tropics. Mol Phylogenet Evol. 35(1):235–253. PMid:15737594. http://dx.doi.org/10.1016/j.ympev.2004.10.004
- Barber PH, Cheng SH, Erdmann MV, Tenggardjaja K, Ambariyanto. 2011. Evolution and conservation of marine biodiversity in the Coral Triangle: insights from stomatopod Crustacea. *In:* Held C, Koenemann S, Schubart CD, editors. Phylogegraphy and Population Genetics in Crustacea. Boca Raton, FL: CRC Press. p. 129–156.
- Barber PH, Erdmann MV, Palumbi SR. 2006. Comparative phylogeography of three codistributed stomatopods: origins and timing of regional lineage diversification in the coral triangle. Evolution. 60(9):1825–1839. PMid:17089967.
- Barber PH, Palumbi SR, Erdmann MV, Moosa MK. 2000. Biogeography—a marine Wallace's line? Nature. 406(6797):692–693. PMid:10963585. http://dx.doi.org/10.1038/35021135
- Barber PH, Palumbi SR, Erdmann MV, Moosa MK. 2002. Sharp genetic breaks among populations of a benthic marine crustacean indicate limited oceanic larval transport: patterns, cause, and consequence. Mol Ecol. 11:659–674. PMid:11972755. http://dx.doi. org/10.1046/j.1365-294X.2002.01468.x
- Barton NH, Hewitt GM. 1985. Analysis of Hybrid Zones. Annu Rev Ecol Syst. 16:113–148. http://dx.doi.org/10.1146/annurev.es.16.110185.000553
- Bellwood DR, Meyer CP. 2009. Endemism and evolution in the Coral Triangle: a call for clarity. J Biogeogr. 36(10):2010–2012. http://dx.doi.org/10.1111/j.1365-2699.2009.02167.x
- Benzie JAH. 1999. Major genetic differences between crown-of-thorns starfish (*Acanthaster planci*) populations in the Indian and Pacific Oceans. Evolution. 53(6):1782–1795. http:// dx.doi.org/10.2307/2640440
- Bird CE, Karl SA, Smouse P, Toonen RJ. 2011. Detecting and measuring and genetic differentiation. In: Held C, Koenemann S, Schubart CD, editors. Phylogegraphy and population genetics in Crustacea. Boca Raton, FL: CRC Press. p. 31–55. http://dx.doi.org/10.1201/b11113-4
- Bremer JRA, Vinas J, Mejuto J, Ely B, Pla C. 2005. Comparative phylogeography of Atlantic bluefin tuna and swordfish: the combined effects of vicariance, secondary contact, introgression, and population expansion on the regional phylogenies of two highly migratory pelagic fishes. Mol Phylogenet Evol. 36(1):169–187. PMid:15904864. http://dx.doi.org/10.1016/j.ympev.2004.12.011
- Briggs JC. 1974. Marine Zoogeography. New York: McGraw-Hill.
- Briggs JC, Bowen BW. 2013. Marine shelf habitat: biogeography and evolution. J Biogeogr. 40:1023-1035. http://dx.doi.org/10.1111/jbi.12082
- Cardinale M, Nugroho D, Hernroth L. 2009. Reconstructing historical trends of small pelagic fish in the Java Sea using standardized commercial trip based catch per unit of effort. Fish Res. 99(3):151–158. http://dx.doi.org/10.1016/j.fishres.2009.05.015
- Carpenter KE, Barber PH, Crandall ED, Ablan-Lagman MCA, Ambariyanto, Mahardika GN, Manjaji-Matsumoto BM, Juinio-Meñez MA, Santos MD, Starger CJ, et al. 2011. Comparative phylogeography of the Coral Triangle and implications for marine management. J Mar Biol. http://dx.doi.org/10.1155/2001/36982.
- Carvalho GR, Hauser L. 1994. Molecular genetics and the stock concept in fisheries. Rev Fish Biol Fish. 4(3):326–350. http://dx.doi.org/10.1007/BF00042908
- Chow S, Ushiama H. 1995. Global population structure of albacore (*Thunnus alalunga*) inferred by RFLP analysis of the mitochondrial ATPase gene. Mar Biol. 123(1):39–45. http:// dx.doi.org/10.1007/BF00350321
- Christie P. 2005. Is integrated coastal management sustainable? Ocean Coastal Manage. 48(3–6):208–232. http://dx.doi.org/10.1016/j.ocecoaman.2005.04.002

- Collette BB. 1995. Scombridae. Atunes, bacoretas, bonitos, caballas, estorninos, melva, etc. *In:* Fischer W, Krupp F, Schneider W, Sommer C, Carpenter KE, Niem V, editors. Guia FAO para Identification de Especies para lo Fines de la Pesca. Pacifico Centro-Oriental. Rome: FAO. p. 1521–1543.
- Collette BB, Nauen CE. 1983. Vol. 2. Scombrids of the world. An annotated and illustrated catalogue of tunas, mackerels, bonitos and related species known to date. Rome: FAO.
- Cook RM, Sinclair A, Stefansson G. 1997. Potential collapse of North Sea cod stocks. Nature. 385(6616):521–522. http://dx.doi.org/10.1038/385521a0
- Crandall ED, Frey MA, Grosberg RK, Barber PH. 2008a. Contrasting demographic history and phylogeographical patterns in two Indo-Pacific gastropods. Mol Ecol. 17(2):611–626. PMid:18179436. http://dx.doi.org/10.1111/j.1365-294X.2007.03600.x
- Crandall ED, Jones ME, Munoz MM, Akinronbi B, Erdmann MV, Barber PH. 2008b. Comparative phylogeography of two seastars and their ectosymbionts within the Coral Triangle. Mol Ecol. 17(24):5276–5290. PMid:19067797. http://dx.doi.org/10.1111/j.1365-294X.2008.03995.x
- DeBoer TS, Subia MD, Ambariyanto, Erdmann MV, Kovitvongsa K, Barber PH. 2008. Phylogeography and limited genetic connectivity in the endangered boring giant clam across the Coral Triangle. Conserv Biol. 22(5):1255–1266. PMid:18637905. http://dx.doi. org/10.1111/j.1523-1739.2008.00983.x
- Diaz-Jaimes P, Uribe-Alcocer M, Rocha-Olivares A, Garcia-de-Leon FJ, Nortmoon P, Durand JD. 2010. Global phylogeography of the dolphin fish (*Coryphaena hippurus*): the influence of large effective population size and recent dispersal on the divergence of marine pelagic cosmopolitan species. Mol Phylogenet Evol. 57:1209–1218. PMid:20971198. http://dx.doi. org/10.1016/j.ympev.2010.10.005
- Drew J, Barber PH. 2009. Sequential cladogenesis of the reef fish *Pomacentrus moluccensis* (Pomacentridae) supports the peripheral origin of marine biodiversity in the Indo-Australian archipelago. Mol Phylogenet Evol. 53(1):335–339. PMid:19401237. http://dx.doi.org/10.1016/j.ympev.2009.04.014
- Drummond AJ, Ashton B, Buxton S, Cheung M, Cooper A, Duran C, Field M, Heled J, Kearse M, Markowitz S, et al. 2011. Geneious. 5.6 ed.
- Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012. Bayesian phylogenetics with BEAUti and the BEAST 1.7. Mol Biol Evol. 29(8):1969–1973. PMid:22367748. PMCid:PMC3408070. http://dx.doi.org/10.1093/molbev/mss075
- Dwipongoo A. 1987. Indonesia's marine fisheries resources. In: Bailey C, Dwiponggo A, Marahudin F, editors. Indonesian marine capture fisheries. Manila, Philippines: International Center for Living Aquatic Resources Management.
- Ekman S. 1953. Zoogeography of the sea. London: Sidgwick and Jackson.
- Ely B, Viñas J, Bremer JRA, Black D, Lucas L, Covelo K, Labrie A, Thelen E. 2005. Consequences of the historical demography on the global population structure of two highly migratory cosmopolitan marine fishes: the yellowfin tuna (*Thunnus albacares*) and the skipjack tuna (*Katsuwonis pelamis*). BMC Evol Biol. 5:19–27. PMid:15725349. PMCid:PMC554763. http://dx.doi.org/10.1186/1471-2148-5-19
- Ersts PJ. Geographic distance matrix generator 1.2.3 ed. American Museum of Natural History, Center for Biodiversity and Conservation. Accessed 19 February 2013. Available from: http://biodiversityinformatics.amnh.org/open_source/gdmg
- Excoffier L, Laval G, Schneider S. 2005. Arlequin (version 3.0): an integrated software package for population genetics data analysis. Evol Bioinform. 1:47–50.
- Excoffier L, Smouse PE, Quattro JM. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes—application to human mitochondrial DNA restriction data. Genetics. 131(2):479–491. PMid:1644282. PMCid:PMC1205020.
- FAO. Fishery and aquaculture country profiles. Indonesia. Rome. Accessed 28 November 2012. Available from: http://www.fao.org/fishery/countrysector/FI-CP_ID/en
- Fleminger A. 1986. The Pleistocene equatorial barrier between the Indian and Pacific oceans and a likely cause for Wallace's Line. Pelagic Biogeogr. 49:84–97.

- Fu YX. 1996. New statistical tests of neutrality for DNA samples from a population. Genetics. 143(1):557–570. PMid:8722804 PMCid:PMC1207287
- Fu YX. 1997. Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. Genetics. 147(2):915–925. PMid:9335623. PMCid:PMC1208208.
- Fu YX, Li WH. 1993. Statistical tests of neutrality of mutations. Genetics. 133(3):693–709. PMid:8454210. PMCid:PMC1205353.
- Gill AC, Allen GR. 2011. *Pseudochromis erdmanni*, a new species of dottyback with medially placed palatine teeth from Indonesia (Teleostei: Perciformes: Pseudochromidae). Zootaxa. 2924:57–62.
- Godfrey JS. 1996. The effect of the Indonesian throughflow on ocean circulation and heat exchange with the atmosphere: a review. J Geophys Res-Oceans. 101(C5):12217–12237. http://dx.doi.org/10.1029/95JC03860
- Gordon AL, Fine RA. 1996. Pathways of water between the Pacific and Indian oceans in the Indonesian seas. Nature. 379(6561):146–149. http://dx.doi.org/10.1038/379146a0
- Graves JE. 1998. Molecular insights into the population structures of cosmopolitan marine fishes. J Hered. 89(5):427–437. http://dx.doi.org/10.1093/jhered/89.5.427
- Graves JE, Ferris SD, Dizon AE. 1984. Close genetic similarity of Atlantic and Pacific skipjack tuna (*Katsuwonus pelamis*) demonstrated with restriction endonuclease analysis of mito-chondrial DNA. Mar Biol. 79(3):315–320. http://dx.doi.org/10.1007/BF00393264
- Hall R. 2002. Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations. J Asian Earth Sci. 20(4):353–431. http://dx.doi.org/10.1016/S1367-9120(01)00069-4
- Hedrick PW. 1999. Perspective: Highly variable loci and their interpretation in evolution and conservation. Evolution. 53(2):313–318. http://dx.doi.org/10.2307/2640768
- Hedrick PW. 2005. A standardized genetic differentiation measure. Evolution. 59(8):1633–1638. PMid:16329237.
- Horne JB, van Herwerden L, Choat JH, Robertson DR. 2008. High population connectivity across the Indo-Pacific: congruent lack of phylogeographic structure in three reef fish congeners. Mol Phylogenet Evol. 49(2):629–638. PMid:18804542. http://dx.doi.org/10.1016/j. ympev.2008.08.023
- Hughes TP, Bellwood DR, Connolly SR. 2002. Biodiversity hotspots, centres of endemicity, and the conservation of coral reefs. Ecol Lett. 5(6):775–784. http://dx.doi. org/10.1046/j.1461-0248.2002.00383.x
- Hutagalung SP. 2012. Export statistics of fishery products 2011. Jakarta, Indonesia: Ministry of Marine Affairs and Fisheries.
- Hutchings JA. 2000. Collapse and recovery of marine fishes. Nature. 406(6798):882–885. PMid:10972288. http://dx.doi.org/10.1038/35022565
- Jones GP, Milicich MJ, Emslie MJ, Lunow C. 1999. Self-recruitment in a coral reef fish population. Nature. 402(6763):802–804. http://dx.doi.org/10.1038/45538
- Jones GP, Planes S, Thorrold SR. 2005. Coral reef fish larvae settle close to home. Curr Biol. 15(14):1314–1318. PMid:16051176. http://dx.doi.org/10.1016/j.cub.2005.06.061
- Klanten OS, Choat JH, van Herwerden L. 2007. Extreme genetic diversity and temporal rather than spatial partitioning in a widely distributed coral reef fish. Mar Biol. 150(4):659–670. http://dx.doi.org/10.1007/s00227-006-0372-7
- Kochzius M, Nuryanto A. 2008. Strong genetic population structure in the boring giant clam, *Tridacna crocea*, across the Indo-Malay Archipelago: implications related to evolutionary processes and connectivity. Mol Ecol. 17(17):3775–3787. PMid:18662232. http://dx.doi. org/10.1111/j.1365-294X.2008.03803.x
- Kool JT, Paris CB, Barber PH, Cowen RK. 2011. Connectivity and the development of population genetic structure in Indo-West Pacific coral reef communities. Glob Ecol Biogeogr. 20(5):695–706. http://dx.doi.org/10.1111/j.1466-8238.2010.00637.x
- Kumar G, Kunal SP, Menezes MR. 2012. Low genetic variation suggest single stock of kawakawa *Euthynnus affinis* (Cantor, 1849) along the Indian Coast. Turk J Fish Aquat Sc. 12:371–380.

Ladd HS. 1960. Origin of the Pacific island molluscan fauna. Am J Sci. 258:137–150.

- Lavery S, Moritz C, Fielder DR. 1996. Indo-Pacific population structure and evolutionary history of the coconut crab *Birgus latro*. Mol Ecol. 5:557–570. http://dx.doi.org/10.1111/j.1365-294X.1996.tb00347.x
- Lee WJ, Conroy J, Howell WH, Kocher TD. 1995. Structure and evolution of teleost mitochondrial control regions. J Mol Evol. 41(1):54–66. PMid:7608989. http://dx.doi.org/10.1007/ BF00174041
- Leray M, Beldade R, Holbrook SJ, Schmitt RJ, Planes S, Bernardi G. 2010. Allopatric divergence and speciation in coral reef fish: the three-spot *Dascyllus, Dascyllus trimaculatus,* species complex. Evolution. 64(5):1218–1230. PMid:20002167.
- Lessios HA, Kessing BD, Pearse JS. 2001. Population structure and speciation in tropical seas: global phylogeography of the sea urchin *Diadema*. Evolution. 55(5):955–975. http://dx.doi. org/10.1554/0014-3820(2001)055[0955:PSASIT]2.0.CO;2
- Librado P, Rozas J. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. Bioinformatics. 25(11):1451–1452. PMid:19346325. http://dx.doi.org/10.1093/bioinformatics/btp187
- Lourie SA, Green DM, Vincent ACJ. 2005. Dispersal, habitat differences, and comparative phylogeography of southeast Asian seahorses (Syngnathidae: *Hippocampus*). Mol Ecol. 14(4):1073–1094. PMid:15773937. http://dx.doi.org/10.1111/j.1365-294X.2005.02464.x
- Lu CP, Chen CA, Hui CF, Tzeng TD, Yeh SY. 2006. Population genetic structure of the swordfish, *Xiphias gladius* (Linnaeus, 1758), in the Indian Ocean and west Pacific inferred from the complete DNA sequence of the mitochondrial control region. Zool Stud. 45(2):269–279.
- Lukas R, Firing E, Hacker P, Richardson PL, Collins CA, Fine R, Gammon R. 1991. Observations of the Mindanao Current during the western equatorial Pacific circulation study. J Geophys Res-Oceans. 96(C4):7089–7104. http://dx.doi.org/10.1029/91JC00062
- Majkowski J. 2007. Global fishery resources of tuna and tuna-like species. Rome, Italy: FAO.
- Manni F, Guerard E, Heyer E. 2004. Geographic patterns of (genetic, morphologic, linguistic) variation: how barriers can be detected by using Monmonier's algorithm. Hum Biol. 76(2):173–190. PMid:15359530. http://dx.doi.org/10.1353/hub.2004.0034
- McCune AR, Lovejoy NR. 1998. The relative rate of sympatric and allopatric speciation in fishes: tests using DNA sequence divergence between sister species and among clades. *In:* Howard DJ, Berlocher SH, editors. Endless forms: species and speciation. New York: Oxford University Press.
- McManus JW. Marine speciation, tectonics and sea-level changes in southeast Asia. 5th International Coral Reef Congress; 1985; Tahiti. p. 133–138.
- McMillan WO, Palumbi SR. 1995. Concordant evolutionary patterns among Indo-West Pacific butterflyfishes. Proc Roy Soc Lond B Bio. 260(1358):229–236. PMid:7784441. http://dx.doi. org/10.1098/rspb.1995.0085
- McQuinn IH. 1997. Metapopulations and the Atlantic herring. Rev Fish Biol Fish. 7(3):297–329. http://dx.doi.org/10.1023/A:1018491828875
- Meirmans PG, Hedrick PW. 2011. Assessing population structure: $F_{\rm ST}$ and related measures. Mol Ecol Resour. 11(1):5–18. PMid:21429096. http://dx.doi. org/10.1111/j.1755-0998.2010.02927.x
- Meirmans PG. 2012. The trouble with isolation by distance. Mol Ecol. 21:2839–2846. PMid:22574758. http://dx.doi.org/10.1111/j.1365-294X.2012.05578.x
- Ministry of Marine Affairs and Fisheries. 2011. Marine and fisheries in figures 2011. Jakarta, Indonesia: Ministry of Marine Affairs and Fisheries. p. 120.
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. Nature. 403(6772):853–858. PMid:10706275. http://dx.doi.org/10.1038/35002501
- Nesbo CL, Rueness EK, Iversen SA, Skagen DW, Jakobsen KS. 2000. Phylogeography and population history of Atlantic mackerel (*Scomber scombrus* L.): a genealogical approach

reveals genetic structuring among the eastern Atlantic stocks. Proc Roy Soc B-Biol Sci. 267(1440):281–292.

- Nuryanto A, Kochzius M. 2009. Highly restricted gene flow and deep evolutionary lineages in the giant clam *Tridacna maxima*. Coral Reefs. 28(3):607–619. http://dx.doi.org/10.1007/s00338-009-0483-y
- Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Henry M, Stevens H, Wagner H. 2013. vegan: community ecology package [computer program]. R package version 2.0-9. Available from: http://cran.r-project.org/web/packages/vegan/index.html
- Palumbi SR, Grabowsky G, Duda T, Geyer L, Tachino N. 1997. Speciation and population genetic structure in tropical Pacific sea urchins. Evolution. 51:1506–1517. http://dx.doi. org/10.2307/2411203
- Pauly D, Cabanban A, Torres FSB. 1996. Fishery biology of 40 trawl-caught teleosts of western Indonesia. *In:* Pauly D, Martosubroto P, editors. Baseline studies of biodiversity: the fish resource of western Indonesia. Makati, Philippines: ICLARM. p. 135–216.
- Perrin C, Borsa P. 2001. Mitochondrial DNA analysis of the geographic structure of Indian scad mackerel in the Indo-Malay Archipelago. J Fish Biol. 59(5):1421–1426. http://dx.doi. org/10.1111/j.1095-8649.2001.tb00205.x
- Planes S, Jones GP, Thorrold SR. 2009. Larval dispersal connects fish populations in a network of marine protected areas. Proc Natl Acad Sci USA. 106(14):5693–5697. PMid:19307588. PMCid:PMC2659712. http://dx.doi.org/10.1073/pnas.0808007106
- Porter SC. 1989. Some geological implications of average Quaternary glacial conditions. Quaternary Res. 32(3):245–261. http://dx.doi.org/10.1016/0033-5894(89)90092-6
- Posada D. 2008. jModelTest: phylogenetic model averaging. Mol Biol Evol. 25(7):1253–1256. PMid:18397919. http://dx.doi.org/10.1093/molbev/msn083
- Ravago-Gotanco RG, Magsino RM, Juinio-Meñez MA. 2007. Influence of the North Equatorial Current on the population genetic structure of *Tridacna crocea* (Mollusca: Tridacnidae) along the eastern Philippine seabord. Mar Ecol Prog Ser. 336:161–168. http://dx.doi. org/10.3354/meps336161
- Reece JS, Bowen BW, Joshi K, Goz V, Larson A. 2010. Phylogeography of two moray eels indicates high dispersal throughout the Indo-Pacific. J Heredity. 57(2):829–835.
- Riede K. 2004. Global register of migratory species from global to regional scales. Bonn, Germany: Federal Agency for Nature Conservation. PMCid:PMC1809154.
- Roberts CM. 1997. Connectivity and management of Caribbean coral reefs. Science. 278(5342):1454–1457. PMid:9367956. http://dx.doi.org/10.1126/science.278.5342.1454
- Roberts CM, McClean CJ, Veron JEN, Hawkins JP, Allen GR, McAllister DE, Mittermeier CG, Schueler FW, Spalding M, Wells F, et al. 2002. Marine biodiversity hotspots and conservation priorities for tropical reefs. Science. 295(5558):1280–1284. PMid:11847338. http:// dx.doi.org/10.1126/science.1067728
- Rohfritsch A, Borsa P. 2005. Genetic structure of Indian scad mackerel *Decapterus russelli:* Pleistocene vicariance and secondary contact in the Central Indo-West Pacific Seas. Heredity. 95(4):315–326. PMid:16094299. http://dx.doi.org/10.1038/sj.hdy.6800727
- Sale PF. 1991. Reef fish communities: open nonequilibrial systems. *In:* Sale PF, editor. The ecology of fishes on coral reefs. San Diego: Academic Press. p. 564-598.
- Santos MD, Lopez GV, Barut NC. 2010. A pilot study on the genetic variation of eastern little tuna (*Euthynnus affinis*) in Southeast Asia. Philipp J Sci. 139(1):43–50.
- Schiller A, Oke PR, Brassington G, Entel M, Fiedler R, Griffin DA, Mansbridge JV. 2008. Eddyresolving ocean circulation in the Asian-Australian region inferred from an ocean reanalysis effort. Prog Oceanogr. 76(3):334–365. http://dx.doi.org/10.1016/j.pocean.2008.01.003
- Springer VG, Williams JT. 1990. Widely distributed Pacific plate endemics and lowered sealevel. Bull Mar Sci. 47(3):631–640.
- Sulaiman ZH, Ovenden JR. 2010. Population genetic evidence for the east-west division of the narrow-barred Spanish mackerel (*Scomberomorus commerson*, Perciformes: Teleostei)

along Wallace's Line. Biodivers Conserv. 19(2):563–574. http://dx.doi.org/10.1007/ s10531-009-9699-y

- Swofford DL. 1993. PAUP a computer program for phylogenetc inference using maximum parsimony. J Gen Physiol. 102(6):A9–A9.
- Tajima F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. Genetics. 123(3):585–595. PMid:2513255. PMCid:PMC1203831.
- Timm J, Figiel M, Kochzius M. 2008. Contrasting patterns in species boundaries and evolution of anemonefishes (Amphiprioninae, Pomacentridae) in the centre of marine biodiversity. Mol Phylogenet Evol. 49(1):268–276. PMid:18513996. http://dx.doi.org/10.1016/j. ympev.2008.04.024
- Utter FM. 1991. Biochemical genetics and fishery management: an historical perspective. J Fish Biol. 39:1–20. http://dx.doi.org/10.1111/j.1095-8649.1991.tb05063.x
- Valsala V, Maksyutov S, Murtugudde R. 2011. Interannual to interdecadal variabilities of the Indonesian Throughflow source water pathways in the Pacific Ocean. J Phys Oceanogr. 41(10):1921–1940. http://dx.doi.org/10.1175/2011JPO4561.1
- van den Bergh GD, de Vos J, Sondaar PY. 2001. The Late Quaternary palaeogeography of mammal evolution in the Indonesian Archipelago. Paleogeogr Palaeocl. 171(3–4):385–408. http://dx.doi.org/10.1016/S0031-0182(01)00255-3
- Vogler C, Benzie JAH, Lessios HA, Barber PH, Worheide G. 2008. A threat to coral reefs multiplied? Four species of crownof-thorns starfish. Biol Lett. 4:696–699. PMid:18832058. PMCid:PMC2614177. http://dx.doi.org/10.1098/rsbl.2008.0454
- Voris HK. 2000. Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. J Biogeogr. 27(5):1153–1167. http://dx.doi. org/10.1046/j.1365-2699.2000.00489.x
- Walsh PS, Metzger DA, Higuchi R. 1991. Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. Biotechniques. 10(4):506–513. PMid:1867860.
- Ward RD, Woodwark M, Skibinski DOF. 1994. A comparison of genetic diversity levels in marine, freshwater, and anadromous fishes. J Fish Biol. 44(2):213–232. http://dx.doi. org/10.1111/j.1095-8649.1994.tb01200.x
- Williams ST, Benzie JAH. 1998. Evidence of a biogeographic break between populations of a high dispersal starfish: congruent regions within the Indo-West Pacific defined by color morphs, mtDNA, and allozyme data. Evolution. 52(1):87–99. http://dx.doi.org/10.2307/2410923
- Woodland DJ. 1983. Zoogeography of the Siganidae (Pisces): an interpretation of distribution and richness patterns. Bull Mar Sci. 33(3):713–717.
- Woodland DJ. 1986. Wallace's Line and the distribution of marine inshore fishes. *In:* Proc Second Int Conf on Indo-Pacific Fishes. Tokyo. p. 453–460. PMid:2426468. PMCid:PMC253096.
- Wray CG, Landman NH, Saunders WB, Bonacum J. 1995. Genetic divergence and geographic diversification in *Nautilus*. Paleobiology. 21(2):220–228.
- Wright S. 1943. Isolation by distance. Genetics. 28(2):114–138. PMid:17247074. PMCid:PMC1209196.
- Wright S. 1951. The genetical structure of populations. Ann Eugenics. 15(4):323-354.
- Wright S. 1965. The interpretation of population structure by *F*-statistics with special regard to systems of mating. Evolution. 19(3):395–420. http://dx.doi.org/10.2307/2406450
- Wyrtki K. 1961. Physical oceanography of the Southeast Asian waters. In: NAGA Report. 15.

